

NHCU-HEP-97-04

NCTU-TH-97-03

UM-HEP-97-10

QCD Corrections to $b \rightarrow s\gamma\gamma$ and Exclusive $B_s \rightarrow \gamma\gamma$ DecayChia-Hung V. Chang ^a, Guey-Lin Lin^{b*} and York-Peng Yao^c^aDepartment of Physics, National Tsing Hua University, Hsinchu, Taiwan, R.O.C.^bInstitute of Physics, National Chiao Tung University, Hsinchu, Taiwan, R.O.C.^cRandall Laboratory of Physics, University of Michigan, Ann Arbor, MI 48109, U.S.A.**Abstract**

The short distance QCD corrections to $b \rightarrow s\gamma\gamma$ are calculated in the leading logarithmic approximation. The equivalence of operator basis reduction for S-matrix elements by using the equations of motion or by proving a low energy theorem is discussed. We apply the above results to the exclusive $B_s \rightarrow \gamma\gamma$ decay. The branching ratio of this decay is found to be 5×10^{-7} in the Standard Model. We also found that QCD corrections modify considerably the ratio between CP-even and CP-odd two-photon amplitudes.

PACS numbers: 13.20.He, 13.40.Hq, 12.38.-t .

Typeset using REVTeX

*E-mail: glin@beauty.phys.nctu.edu.tw, Tel: 886-3-5731984, Fax: 886-3-5720728.

Rare B decays provide useful probes to the flavor structures of the Standard Model(SM) and its various extensions. Among them, the radiative decay $B \rightarrow X_s \gamma$ [1] has recently received the most attention since it is not only within the reach of current experiments, where CLEO [2] gives an updated branching ratio

$$Br(B \rightarrow X_s \gamma) = (2.32 \pm 0.51 \pm 0.29 \pm 0.32) \times 10^{-4}. \quad (1)$$

but also theoretically it has been calculated to the NLO accuracy [3,4] with the result

$$Br(B \rightarrow X_s \gamma) = (3.28 \pm 0.33) \times 10^{-4}, \quad (2)$$

in the SM. While this decay mode will be continually studied for some time, the upcoming B factories at KEK, SLAC and hadronic B projects at HERA, Tevatron and LHC offer new opportunities to study many more rare decay modes. With these facilities, it will be possible to study decay modes with branching ratios as small as 10^{-8} . In this work, we focus on the exclusive decay $B_s \rightarrow \gamma\gamma$, which, in spite of its small branching ratio, has a very clear signal where two monochromatic energetic photons are produced precisely back-to-back in the rest frame of B_s . Within the SM, the electroweak contributions to this process was calculated without QCD corrections some time ago [5–7]. The branching ratio so obtained was about 10^{-7} . The calculation was later extended [8] to the case of two-Higgs doublet models(2HDM). It was shown that $Br(B_s \rightarrow \gamma\gamma)$ might be enhanced to 10^{-6} in certain regions of the parameter space. The current experimental bound on this decay is $Br(B_s \rightarrow \gamma\gamma) < 1.48 \times 10^{-4}$ [9].

Before one goes on to study other *new physics* which potentially can influence this decay, it stands to reason to improve upon previous calculations by incorporating short distance QCD running from the scale m_W to the scale m_b . This will be achieved by constructing an effective Hamiltonian for the quark level process $b \rightarrow s\gamma\gamma$ at the scale $\mu = m_b$. The amplitude for $B_s \rightarrow \gamma\gamma$ can then be determined once a method is devised for computing the hadronic matrix elements.

At present, the effective Hamiltonian $H_{eff}(b \rightarrow s\gamma)$ for $b \rightarrow s\gamma$ is already well understood [10]. It is natural to ask what is the corresponding effective Hamiltonian for $b \rightarrow s\gamma\gamma$. Is it

identical to $H_{eff}(b \rightarrow s\gamma)$ or more complicated with extra operators? The answer is that, to the order $1/m_W^2$, $H_{eff}(b \rightarrow s\gamma)$ is sufficient to generate all contributions to $b \rightarrow s\gamma\gamma$. This can be understood by either applying the equations of motion [12] or by applying an extension of Low's low energy theorem [5,13]. In the following we shall illustrate that the above two approaches are in fact equivalent. To proceed, we write the effective Hamiltonian for $b \rightarrow s\gamma$ as [1]

$$H_{eff}(b \rightarrow s\gamma) = -\frac{G_F}{\sqrt{2}} V_{ts}^* V_{tb} \sum_{i=1}^8 C_i(\mu) O_i(\mu), \quad (3)$$

where

$$\begin{aligned} O_1 &= (\bar{s}_i c_j)_{V-A} (\bar{c}_j b_i)_{V-A} \\ O_2 &= (\bar{s}_i c_i)_{V-A} (\bar{c}_j b_j)_{V-A} \\ O_3 &= (\bar{s}_i b_i)_{V-A} \sum_q (\bar{q}_j q_j)_{V-A} \\ O_4 &= (\bar{s}_i b_j)_{V-A} \sum_q (\bar{q}_j q_i)_{V-A} \\ O_5 &= (\bar{s}_i b_i)_{V-A} \sum_q (\bar{q}_j q_j)_{V+A} \\ O_6 &= (\bar{s}_i b_j)_{V-A} \sum_q (\bar{q}_j q_i)_{V+A} \\ O_7 &= \frac{e}{4\pi^2} \bar{s}_i \sigma^{\mu\nu} (m_s P_L + m_b P_R) b_i F_{\mu\nu} \\ O_8 &= \frac{g}{4\pi^2} \bar{s}_i \sigma^{\mu\nu} (m_s P_L + m_b P_R) T_{ij}^a b_j G_{\mu\nu}^a. \end{aligned} \quad (4)$$

We remark that the decay $b \rightarrow s\gamma\gamma$ also receives contributions induced by γ and Z^0 penguin diagrams, which are higher-order in electroweak couplings. Effects of such diagrams can be summarized into electroweak penguin operators [11] which have similar structures as strong penguins $O_3 - O_6$, and an $O(\alpha_{em})$ -correction to the Wilson coefficient C_3 . It is well known that [11] Z^0 penguin diagram contains the $O(m_t^2)$ -contribution which is not much suppressed compared to contributions of strong penguins. On the other hand, as we shall see later, the dominant contributions to $b \rightarrow s\gamma\gamma$ actually arise from O_1 , O_2 and O_7 . Therefore, in view of simplifying our analysis, we shall neglect electroweak penguin contributions.

Concerning the effective Hamiltonian $H_{eff}(b \rightarrow s\gamma)$, we note that operator mixings

between the set, $O_1, \dots, O_6$, and the magnetic penguin O_7, O_8 are generally regularization-scheme dependent. However, as pointed out by Ciuchini *et al.* in Ref. [1], the one-loop matrix elements of both $b \rightarrow s\gamma$ and $b \rightarrow sg$ arising from $O_1, \dots, O_6$ carry compensating scheme-dependence such that the total physical amplitudes are independent of regularization schemes. In view of this, one defines the so called “effective Wilson coefficients” [14], $C_7^{eff}(\mu)$ and $C_8^{eff}(\mu)$, for operators O_7 and O_8 , for which the RG running is scheme independent. In ’t Hooft-Veltman’s (HV) scheme for γ_5 , these effective coefficients coincide with original ones. Furthermore, in this scheme, only O_7 contributes to the one-loop matrix element of $b \rightarrow s\gamma$ [15]. Since we shall adopt the HV scheme for subsequent calculations, we drop the superscript “eff” on both C_7 and C_8 from now on.

To see whether $O_1, \dots, O_7$ are sufficient to generate all contributions to $b \rightarrow s\gamma\gamma$, one notes that operators containing two photon fields are missing from Eq. (4). Photon fields which constitute operators in this category can come in 3 ways: from covariant derivatives D_μ , from field strength tensors $F_{\mu\nu}$ or from some combinations of the two. It has been shown by Grinstein *et al.* in Ref. [1] that such operators can be eliminated or simply reduced to those in Eq. (4) by applying equations of motion. Therefore, $H_{eff}(b \rightarrow s\gamma)$ is also the effective Hamiltonian for $b \rightarrow s\gamma\gamma$. At this point, we wish to emphasize that the above simplifications by using the equations of motion are applicable only with respect to the S matrix elements [12]. The Green functions depend on the eliminated operators, however.

The above procedure based on applying the equations of motion has a close link with the low energy theorem derived by us some time ago [5]. Let us illustrate this with the operator $Q_4 \equiv \bar{s}_L \not{D}^3 b_L$ [16]. Upon applying the equation of motion for the b quark, it is easily seen that $Q_4 = im_b^3 \bar{s}_L b_R$. In this new form, Q_4 gives rise to the flavor changing self energy. However this contribution will be subtracted out by the on-shell renormalization [17]

$$\bar{s}\Sigma(P^2 = m_s^2) = 0, \quad \Sigma(P^2 = m_b^2)b = 0. \quad (5)$$

Hence, with equations of motion one concludes that Q_4 gives no contributions to $b \rightarrow s\gamma\gamma$.

This fact is also realized by the low energy theorem. There one takes Q_4 as it is initially defined and computes Q_4 's contributions to $b - s$ self energy, 3-point $b \rightarrow s\gamma$ and 4-point $b \rightarrow s\gamma\gamma$ vertices. Its total contributions to $b \rightarrow s\gamma\gamma$ are given by summing up the irreducible and reducible diagrams where the latter come from attaching *bremsstrahlung* photons to the flavor changing self-energy and the three-point $b \rightarrow s\gamma$ vertex. The renormalization condition, Eq. (5), ensures that the end result will be local, and one can further show that the extraneous contributions cancel out due to gauge invariance [5]. Such a cancellation among reducible and irreducible digrams is what was referred to as the generalization of Low's low energy theorem [5,13]. It is clear that this theorem is essentially a verification of general arguments based upon applying the equations of motion.

With the above issues clarified, we are ready to compute $b \rightarrow s\gamma\gamma$. The amplitude can be separated into irreducible and reducible parts, as shown in Figs. 1 and 2.

Figure 1

Figure 2

We begin with irreducible diagrams which are more involved. There is only one type of diagrams which are depicted generically by Figure 1, where the four-point vertex is a result of inserting 4-quark operators $O_1 - O_6$. While in the case of O_1 and O_2 , Figure 1 consists of only one diagram, there are, however, two different ways of contracting in the case of $O_3 - O_6$, as the light quark field q appearing in these operators can be external particles as well. To simplify the algebra, we apply Fierz rearrangements to four-quark operators when necessary. This rearrangement is legitimate because the γ_5 we use is simply the product of $\gamma^0, \dots, \gamma^3$. The Fierz identities we have used are

$$\begin{aligned}
(\bar{q}_1 q_2)_{V-A} (\bar{q}_3 q_4)_{V-A} &= (\bar{q}_1 q_4)_{V-A} (\bar{q}_3 q_2)_{V-A}, \\
(\bar{q}_1 q_2)_{V-A} (\bar{q}_3 q_4)_{V+A} &= -2 (\bar{q}_1 (1 + \gamma_5) q_4) (\bar{q}_3 (1 - \gamma_5) q_2).
\end{aligned} \tag{6}$$

The amplitude for the irreducible diagrams can be written as

$$\begin{aligned}
M_{IR} &= \frac{16\sqrt{2}\alpha G_F}{9\pi} V_{ts}^* V_{tb} \times \\
&\bar{u}(p') \left[\sum_q A_q J(m_q^2) \gamma^\rho P_L R_{\mu\nu\rho} + iB \left(m_s K(m_s^2) P_L + m_b K(m_b^2) P_R \right) T_{\mu\nu} \right. \\
&\left. + C \left(-m_s L(m_s^2) P_L + m_b L(m_b^2) P_R \right) \varepsilon_{\mu\nu\alpha\beta} k_1^\alpha k_2^\beta \right] u(p) \epsilon^\mu(k_1) \epsilon^\nu(k_2),
\end{aligned} \tag{7}$$

where

$$\begin{aligned}
R_{\mu\nu\rho} &= k_{1,\nu} \varepsilon_{\mu\rho\sigma\lambda} k_1^\sigma k_2^\lambda - k_{2,\mu} \varepsilon_{\nu\rho\sigma\lambda} k_1^\sigma k_2^\lambda \\
&+ k_1 \cdot k_2 \varepsilon_{\mu\nu\rho\sigma} (k_2 - k_1)^\sigma
\end{aligned} \tag{8}$$

is a third rank tensor originally constructed by Rosenberg and Adler [18]; and

$$T_{\mu\nu} = (k_{2,\mu} k_{1,\nu} - g_{\mu\nu} k_1 \cdot k_2). \tag{9}$$

Here we adopt the convention: $\varepsilon_{0123} = 1$. The coefficients A_q 's, B and C are linear combinations of Wilson coefficients and are given by

$$\begin{aligned}
A_u &= (C_3 - C_5) N_c + (C_4 - C_6) \\
A_d &= \frac{1}{4} ((C_3 - C_5) N_c + (C_4 - C_6)) \\
A_c &= ((C_1 + C_3 - C_5) N_c + (C_2 + C_4 - C_6)) \\
A_s &= A_b = \frac{1}{4} ((C_3 + C_4 - C_5) N_c + (C_3 + C_4 - C_6)) \\
B &= C = -\frac{1}{4} (C_6 N_c + C_5).
\end{aligned} \tag{10}$$

Note that the above Wilson coefficients are evaluated at scale $\mu = m_b$. Finally the functions $J(m^2)$, $K(m^2)$ and $L(m^2)$ are defined by

$$\begin{aligned}
J(m^2) &= I_{11}(m^2), \\
K(m^2) &= \left(4I_{11}(m^2) - I_{00}(m^2) \right), \\
L(m^2) &= I_{00}(m^2),
\end{aligned} \tag{11}$$

with

$$I_{pq}(m^2) = \int_0^1 dx \int_0^{1-x} dy \frac{x^p y^q}{m^2 - 2xy k_1 \cdot k_2 - i\varepsilon}. \quad (12)$$

For the reducible diagrams, the one loop contributions with O_1 – O_6 insertion actually vanish, similar to what occurs in $b \rightarrow s\gamma$ in HV-scheme. Therefore the amplitude for reducible diagrams is identical to its lowest order form [5] except for replacing $C_7(m_W)$ by $C_7(m_b)$. Hence

$$\begin{aligned} M_R = & \frac{i\sqrt{2}\alpha G_F}{6\pi} V_{ts}^* V_{tb} C_7(m_b) \times \\ & \bar{u}(p') \left[\left(\frac{1}{p' \cdot k_1} - \frac{1}{p \cdot k_2} \right) \sigma_{\mu\beta} \sigma_{\nu\alpha} k_1^\beta k_2^\alpha + 2i \left(\frac{p'_\mu}{p' \cdot k_1} - \frac{p_\mu}{p \cdot k_1} \right) \sigma_{\nu\beta} k_2^\beta \right] \\ & \cdot (m_b P_R + m_s P_L) u(p) \epsilon^\mu(k_1) \epsilon^\nu(k_2) + k_1, \mu \longleftrightarrow k_2, \nu. \end{aligned} \quad (13)$$

From Eqs. (7) and (13), we obtain the total amplitude for $b \rightarrow s\gamma\gamma$.

To calculate $B_s \rightarrow \gamma\gamma$, one may follow a perturbative QCD approach which includes a proof of factorization, showing that soft gluon effects can be factorized into B_s meson wave function; and a systematic way of resumming large logarithms due to “hard gluons” with energies between 1 GeV and m_b . Since such an approach is technically rather demanding with uncertain reliability, we shall adapt a phenomenological approach where the long distance effects are replaced by a few non-perturbative parameters. In other words we simply evaluate the hadronic matrix element of $M_{IR} + M_R$, relying on a phenomenological model. Before doing so, it is important to note that M_R is apparently non-local due to internal b or s quark propagators. To handle these non-local terms, one observes that the b quark inside the B_s meson carries most of the meson energy, and its four velocity can be treated as equal to that of B_s . Hence one may write b quark momentum as $p = m_b v$ where v is the common four velocity of b and B_s . With this parametrization, we have

$$\begin{aligned} p \cdot k_1 &= m_b v \cdot k_1 = \frac{1}{2} m_b m_{B_s} = p \cdot k_2, \\ p' \cdot k_1 &= (p - k_1 - k_2) \cdot k_1 = -\frac{1}{2} m_{B_s} (m_{B_s} - m_b) = p' \cdot k_2, \end{aligned} \quad (14)$$

where the second equation is based on a constituent picture [5] that b and $\bar{s}$ quarks share the

total energy of B_s ¹. Therefore $-p'$ should be taken as the four momentum of a constituent $\bar{s}$ quark. With Eq. (14), M_R is readily made local. We then compute the amplitude for $B_s \rightarrow \gamma\gamma$ using the following relations

$$\begin{aligned}\langle 0|\bar{s}\gamma_\mu\gamma_5 b|B_s(P)\rangle &= -if_{B_s}P_\mu, \\ \langle 0|\bar{s}\gamma_5 b|B_s(P)\rangle &= if_{B_s}M_B,\end{aligned}\tag{15}$$

where f_{B_s} is the B_s meson decay constant which is about 200 MeV according to recent Lattice QCD calculations [19].

The total amplitude is now separated into a CP-even and a CP-odd part

$$T(B_s \rightarrow \gamma\gamma) = M^+ F_{\mu\nu} F^{\mu\nu} + iM^- F_{\mu\nu} \tilde{F}^{\mu\nu}.\tag{16}$$

We find that

$$M^+ = -\frac{4\sqrt{2}\alpha G_F}{9\pi} f_{B_s} m_{B_s} V_{ts}^* V_{tb} \left(Bm_b K(m_b^2) + \frac{3C_7}{8\bar{\Lambda}} \right),\tag{17}$$

and

$$M^- = \frac{4\sqrt{2}\alpha G_F}{9\pi} f_{B_s} m_{B_s} V_{ts}^* V_{tb} \left(\sum_q m_{B_s} A_q J(m_q^2) + m_b BL(m_b^2) + \frac{3C_7}{8\bar{\Lambda}} \right),\tag{18}$$

where $\bar{\Lambda} = m_{B_s} - m_b$. The decay width for $B_s \rightarrow \gamma\gamma$ is simply

$$\Gamma(B_s \rightarrow \gamma\gamma) = \frac{m_{B_s}^3}{16\pi} (|M^+|^2 + |M^-|^2).\tag{19}$$

To obtain numerical results, we have set light quark masses to zero and used [20] $m_t = 175$ GeV, $m_b = 4.8$ GeV and $m_c = 1.5$ GeV. Furthermore, we take $m_{B_s} = 5.37$ GeV, $\alpha = \frac{1}{129}$ and $V_{ts}^* V_{tb} = 4 \times 10^{-2}$. The numerical values for Wilson coefficients $C_1 - C_8$ evaluated at $\mu = m_b$ are listed in Table I. With the above input parameters, we find $\Gamma(B_s \rightarrow \gamma\gamma) = 2.0 \times 10^{-10}$ eV which amounts to a branching ratio $Br(B_s \rightarrow \gamma\gamma) = 5.0 \times 10^{-7}$, for the given $\Gamma_{B_s}^{total} = 4 \times 10^{-4}$ eV. If QCD corrections are not included, namely taking

¹Note that the momentum of $\bar{s}$ quark is $-p'$ as p' denotes the momentum of s quark in $b \rightarrow s\gamma\gamma$.

$C_2 = C_2(m_W) = 1$, $C_7 = C_7(m_W)$ and setting all the other Wilson coefficients to zero, one obtains $\Gamma(B_s \rightarrow \gamma\gamma) = 1.3 \times 10^{-10}$ eV. Therefore including QCD effects has increased the rate of $B_s \rightarrow \gamma\gamma$ by more than 50%.

It is interesting to note that QCD correction modifies contributions of irreducible and reducible diagrams in opposite ways. It enhances the contributions of reducible diagrams through the enhancement of C_7 , namely $C_7(m_b)/C_7(m_W) \approx 1.5$. Such an enhancement is already well known in the decay $b \rightarrow s\gamma$. As a contradistinction, QCD suppresses the contributions of irreducible diagrams since its effect essentially replaces $C_2(m_W)$ with $N_c C_1(m_b) + C_2(m_b)^2$. The latter is much smaller than the former due to cancellations between $N_c C_1(m_b)$ and $C_2(m_b)$. Because QCD gives distinct effects to M_{IR} and M_R , the relative magnitude of M^+ and M^- is also modified accordingly and drastically by QCD. From Eqs. (10), (17), (18) and Table 1, it is clear that the magnitudes of M^+ and M^- are almost identical, both dominated by C_7 but somewhat corrected by the suppressed $N_c C_1(m_b) + C_2(m_b)$. Numerically we have $|M^+|^2/|M^-|^2 = 0.80$. The corresponding ratio without QCD corrections is 0.38 which is twice smaller.

On the experimental side, as mentioned before, the future B factories are capable of probing B decays with branching ratio as low as 10^{-8} . Therefore one expects the decay $B_s \rightarrow \gamma\gamma$ to be seen in these future facilities. It is, however, more challenging to separate partial amplitudes M^+ and M^- because it will require measuring the spin correlation of final state photons.

At this point, we wish to state the obvious interdependence between $B_s \rightarrow \gamma\gamma$ and inclusive $B \rightarrow X_s \gamma$ decays. It is well known that the later decay depends on $C_7(m_W)$ at the tree level, and $C_8(m_W)$ and $C_2(m_W)$ through QCD-induced operator mixings [1]. CLEO [2] measures some combinations of these coefficients, and thus imposes constraints [14,21]

²Since $C_3(m_b), \dots, C_6(m_b)$ are much smaller than $C_1(m_b)$ and $C_2(m_b)$, the dominant contribution to M_{IR} is proportional to A_c which is approximately equal to $N_c C_1(m_b) + C_2(m_b)$.

on various extensions of the Standard Model, such as 2HDM or Minimal Supersymmetric Standard Model(MSSM). Since $B_s \rightarrow \gamma\gamma$ depends on the same set of Wilson coefficients, its sensitivity to physics beyond the Standard Model complements the corresponding sensitivity in $B \rightarrow X_s\gamma$ [22].

Before closing, we like to comment on the inclusive $B \rightarrow X_s\gamma\gamma$ decay. This process has recently been explored without the inclusion of QCD corrections [23]. The branching ratio is found to be 10^{-7} in the SM. Experimentally, due to the appearance of hadronic final state, it seems to be more difficult to analyze this process than one can do for $B_s \rightarrow \gamma\gamma$. Besides, photons in this decay may be soft or collinear which may require extra treatment such as the necessity of imposing kinematical cuts. On the theoretical side, there exists a further complication in that photons can be emitted from the spectator quark which forms B with b . Such spectator effects, while neglected in Ref. [23], need to be studied carefully in order that additiona

l informations on C_2 and C_7 can be extracted from $B \rightarrow X_s\gamma\gamma$.

In conclusion, we have outlined the equivalence of procedure by using the equations of motion [12] and that by the low energy theorem [5,13] to reduce the operator basis, as they are applied to $b \rightarrow s\gamma\gamma$. We have also computed the exclusive decay $B_s \rightarrow \gamma\gamma$ with the effective Hamiltonian $H_{eff}(b \rightarrow s\gamma)$. The branching ratio is found to be 5×10^{-7} which is 50% larger than the result without QCD corrections. Finally we have argued that one should include spectator effects in the study of inclusive $B \rightarrow X_s\gamma\gamma$ decay.

Note added. After completing this work, we became aware of a paper by G. Hiller and E. O. Ilhan, hep-ph/9704385, where the QCD correction to $B_s \rightarrow \gamma\gamma$ is also discussed. However, the crucial contribution from O_1 and other contributions from $O_3, \dots, O_6$ are not taken into account in that paper.

ACKNOWLEDGMENTS

We thank R. Akhoury for a discussion, and L. Reina for pointing out a sign error and a misprint in the earlier version of the manuscript. GLL also likes to thank particle theory group at U. of Michigan for their hospitality, where part of this work is done. The works of CHC and GLL are supported in part by National Science Council of R.O.C. under grant numbers NSC 86-2112-M-007-020 and NSC 86-2112-M-009-012; YPY's work is partially supported by the U. S. Department of Energy.

REFERENCES

- [1] B. Grinstein, R. Springer and M. B. Wise, Phys. Lett. **B 202** (1988) 138; Nucl. Phys. **B339** (1990) 269; R. Grigjanis, P. J. O' Donnell, M. Sutherland and H. Navelet, Phys. Lett. **B 213** (1988) 355; *ibid.* **B 286** (1992) E, 413; M. Misiak, Phys. Lett. **B 269** (1991) 161; K. Adel and Y.-P. Yao, Mod. Phys. Lett. **A 8** (1993) 1679; M. Ciuchini, E. Franco, G. Martinelli, L. Reina and L. Silvestrini, Phys. Lett. **B 316** (1993) 127; Nucl. Phys. **B 421** (1994) 41; G. Cella, G. Curci, G. Ricciardi and A. Viceré, Phys. Lett. **B 325** (1994) 227; Nucl. Phys. **B 431** (1994) 417; M. Misiak, Nucl. Phys. **B 393** (1993) 23; *ibid.* **B 439** (1995) E, 461.
- [2] CLEO Collaboration R. Ammar *et al.*, Phys. Rev. Lett. **71** (1993) 674; M. S. Alam *et al.*, *ibid.* **74** (1995) 2885.
- [3] K. Adel and Y.-P. Yao, Phys. Rev. **D 49** (1994) 4945; C. Greub and T. Hurth, hep-ph/9703349.
- [4] K. G. Chetyrkin, M. Misiak and M. Münz, hep-ph/9612313.
- [5] G.-L. Lin, J. Liu and Y.-P. Yao, Phys. Rev. Lett. **64** (1990) 1498; Phys. Rev. **D 42** (1990) 2314.
- [6] H. Simma and D. Wyler, Nucl. Phys. **B 344** (1990) 283.
- [7] S. Herrlich and J. Kalinowski, Nucl. Phys. **B 381** (1992) 501.
- [8] T. M. Aliev and G. Turan, Phys. Rev. **D 48** (1993) 1176.
- [9] M. Acciarri *et al.* (L3 Collaboration), Phys. Lett. **B 363** (1995) 127.
- [10] For a review on earlier literatures, see G. Buchalla, A. J. Buras and M. E. Lautenbacher, Review of Modern Physics, **68** (1996) 1125.
- [11] J. M. Flynn and L. Randall, Phys. Lett. **B 224** (1989) 221; A. J. Buras, M. Jamin, M. E. Lautenbacher and P. Weisz, Nucl. Phys. **B 400** (1993) 37; A. J. Buras, M. Jamin

- and M. E. Lautenbacher, *ibid.* **B 400** (1993) 75; M. Ciuchini, E. Franco, G. Martinelli and L. Reina, Nucl. Phys. **B 415** (1994) 403.
- [12] General discussions on the uses of equations of motion in field theory can be found in H. D. Politzer, Nucl. Phys. **B 172** (1980) 349; C. Arzt, Phys. Lett. **B 342** (1995) 189. For specific discussions with respect to rare B decays, see H. Simma, Z. Phys. **C 61** (1994) 67.
- [13] F. E. Low, Phys. Rev. **110** (1958) 974.
- [14] A. J. Buras, M. Misiak, M. Münz and S. Pokorski, Nucl. Phys. **B 424** (1994) 374.
- [15] See discussions by Ciuchini *et al.* in Ref. [1].
- [16] This labeling is according to the convention of Grinstein *et al.* in Ref. [1].
- [17] N. G. Deshpande and G. Eilam, Phys. Rev. **D 26** (1982) 2463.
- [18] L. Rosenberg, Phys. Rev. **129** (1963) 2786; S. L. Adler, *ibid.* **177** (1969) 2426.
- [19] For a review of recent progress, see J. Flynn, in International Conference on High Energy Physics, Warsaw, Poland, 1996.
- [20] Particle Data Group, R. M. Barnett *et al.* Phys. Rev. **D 54** (1996) 1.
- [21] R. Barbieri and G. F. Giudice, Phys. Lett. **B 309** (1993) 86; R. Garisto and J. N. Ng, *ibid.* **B 315** (1993) 372.
- [22] C.-H. V. Chang, G.-L. Lin and Y.-P. Yao, work in progress.
- [23] L. Reina, G. Ricciardi and A. Soni, Phys. Lett. **B 396** (1997) 231.

FIGURES

FIG. 1. The generic irreducible diagram contributing to $b \rightarrow s\gamma\gamma$. The cross in a circle denotes insertions of operators, $O_1, \dots, O_6$. We omit the diagram with two photon lines interchanged.

FIG. 2. The reducible diagrams contributing to $b \rightarrow s\gamma\gamma$ where the interchange of two photon lines is assumed. The cross in a circle denotes the insertion of operator O_7 .

TABLES

TABLE I. Wilson coefficients $C_i(\mu)$ at $\mu = m_b = 4.8$ GeV in the leading logarithmic approximations. These values are obtained by taking $M_W = 80.2$ GeV and $\alpha_s(M_Z) = 0.117$

$C_1(\mu)$	$C_2(\mu)$	$C_3(\mu)$	$C_4(\mu)$	$C_5(\mu)$	$C_6(\mu)$	$C_7(\mu)$	$C_8(\mu)$
-0.222	1.09	0.010	-0.023	0.007	-0.028	-0.301	-0.144




Figure 1




Figure 2